

A vector bundle proof of Poncelet theorem

Jean Vallès*

Université de Pau et des Pays de l'Adour
Laboratoire de Mathématique- Bât. IPRA

Avenue de l'Université
64000 PAU (FRANCE)

e-mail : jean.valles@univ-pau.fr

tel : +33(0)559407517

MSC 2010 Subject Classifications. Primary 14C20, 14J60;
secondary 14M12, 14H50

*Author partially supported by ANR-09-JCJC-0097-0 INTERLOW and ANR GEOLMI

Abstract

There are few different proofs of the celebrated Poncelet closure theorem about polygons simultaneously inscribed in a smooth conic and circumscribed around another. We propose a new proof, based on the link between Schwarzenberger bundles and Poncelet curves.

Keywords: Poncelet porism, Poncelet curves, Darboux theorem, Schwarzenberger bundles.

1 Introduction

In the town of Saratov where he was prisoner, Poncelet, continuing the work of Chapple on triangles simultaneously inscribed in a circle and circumscribed around another circle, proved the following generalization.

Theorem *Let C and D be two smooth conics in $\mathbb{P}^2(\mathbb{C})$ such that there exists one n -gon (polygon with n sides) inscribed in D and circumscribed around C . Then there are infinitely many of such n -gons.*

According to Berger ([2], page 256) this theorem is the nicest result about the geometry of conics (¹). Even if it is, there are few proofs of it. To my knowledge there are only three. The first proof, published in 1822 and based on infinitesimal deformations, is due to Poncelet ([7], page 362). Later, Jacobi proposed a new proof based on finite order points on elliptic curves; his proof, certainly the most famous, is explained in a modern way and in detail by Griffiths and Harris (see [5]). In 1870 Weyr proved a Poncelet theorem in space (more precisely for two quadrics) that implies the one above when one quadric is a cone; this proof is explained by Barth and Bauer ([1], thm. 1.1).

Our aim in this short note is to involve vector bundles techniques to propose a new proof of this celebrated result. Poncelet did not appreciate Jacobi's for the reason that it was too far from the geometric intuition. I guess that he would not appreciate our proof either for the same reason.

2 Preliminaries

In all this text the ground field is \mathbb{C} . A n -gon consists in the data of n distinct ordered points a_1, \dots, a_n and n lines $(a_1a_2), \dots, (a_{n-1}a_n), (a_na_1)$. A complete n -gon consists in the data of n distinct lines in linear general position and their $\binom{n}{2}$ points of intersection, also called vertices.

We say that a n -gon (respectively a complete n -gon) is inscribed in a given curve if this curve passes through its n points (respectively its $\binom{n}{2}$ vertices). We say that a n -gon, or a complete n -gon, is circumscribed around a smooth conic C if the sides of the polygon, i.e. the n lines, are tangent to C .

¹As suggested by the referee, we could name also, among the nicest theorems about conics, the one, due to Chasles, which says that there are 3264 conics tangent to five given conics.




Figure 1: A 17-gon simultaneously inscribed in a conic and circumscribed around another.

3 Schwarzenberger bundles

First of all let us introduce a vector bundle $E_{n,C}$ naturally associated to any set of n lines tangent to a fixed smooth conic $C \subset \mathbb{P}^2$ (they were defined by Schwarzenberger in [8]). A set of n lines tangent to C corresponds by projective duality to a degree n divisor on the dual conic $C^\vee \subset \mathbb{P}^{2\vee}$. According to the isomorphism $C^\vee \simeq \mathbb{P}^1$ we can consider the subvariety $X \subset \mathbb{P}^2 \times \mathbb{P}^1$ defined by the equation $x_0u^2 + x_1uv + x_2v^2 = 0$ ((x_0, x_1, x_2) are the homogeneous coordinates on \mathbb{P}^2 and (u, v) the homogeneous coordinates on \mathbb{P}^1) and the projections p and q respectively on \mathbb{P}^2 and \mathbb{P}^1 .

$$\begin{array}{ccc} X & \xrightarrow{q} & \mathbb{P}^1 \\ p \downarrow & & \\ \mathbb{P}^2 & & \end{array}$$

The variety X is the incidence variety of couples (x, l) such that $x \in \mathbb{P}^2$, l is tangent to C and $x \in l$. It is a double cover of \mathbb{P}^2 ramified along the curve C defined by the vanishing of the discriminant $x_1^2 - 4x_0x_2 = 0$ of the equation of X in $\mathbb{P}^2 \times \mathbb{P}^1$. If $z \in \mathbb{P}^1$ then $p(q^{-1}(z))$ is a line in \mathbb{P}^2 tangent to C . If, instead of considering a point, we are considering a line bundle on \mathbb{P}^1 we will find a vector bundle of rank two on \mathbb{P}^2 by taking the direct image of its inverse image. Moreover, following Schwarzenberger, we know a very explicit resolution of this bundle. Indeed, tensoring the following exact sequence

$$0 \longrightarrow \mathcal{O}_{\mathbb{P}^2 \times \mathbb{P}^1}(-1, -2) \longrightarrow \mathcal{O}_{\mathbb{P}^2 \times \mathbb{P}^1} \longrightarrow \mathcal{O}_X \longrightarrow 0,$$

Remark 4.2. Since the vector space of curves of degree $n - 2$ has dimension $\binom{n}{2}$, the degree of a curve passing through the $\binom{n}{2}$ vertices of a complete n -gon is at least equal to $n - 1$.

Proof. I recall here a proof already written in [9]. A complete n -gon circumscribed around C and inscribed into S corresponds to a non-zero global section $s \in H^0(E_{n,C})$ vanishing along its vertices $Z(s)$:

$$0 \longrightarrow \mathcal{O}_{\mathbb{P}^2} \longrightarrow E_{n,C} \longrightarrow \mathcal{I}_{Z(s)}(n-1) \longrightarrow 0.$$

By the remark of the previous section (2), the set $Z(s)$ belongs to S . It implies that the curve S corresponds to a global section of $\mathcal{I}_{Z(s)}(n-1)$. Since the map

$$H^0(E_{n,C}) \longrightarrow H^0(\mathcal{I}_{Z(s)}(n-1))$$

is surjective, this section comes from a non-zero section $t \in H^0(E_{n,C})$ (i.e. another n -gon) such that $Z(t)$ belongs to S . We deduce then that the determinant

$$\mathcal{O}_{\mathbb{P}^2}^2 \xrightarrow{(s,t)} E_{n,C}$$

is the equation of S . In other terms, a general section in the pencil generated by s and t (i.e a general linear combination of s and t) corresponds to a complete n -gon circumscribed around C and inscribed in S . This proves the theorem. \square




Figure 2: A complete circumscribed 4 gon and a cubic Poncelet curve.

These curves described by Darboux are called *Poncelet curves*. When $n = 5$ they are the so-called Luröth quartics (see [6]).

5 Poncelet theorem

Let us now consider n -gons that are simultaneously inscribed in a smooth conic and circumscribed around another one. For these configurations Poncelet obtained his famous result ([7], page 362). We prove it now, with the help of Poncelet curves defined above.

Theorem 5.1. *Let $C \subset \mathbb{P}^2$ and $D \subset \mathbb{P}^2$ be two smooth conics such that there exists one n -gon inscribed in D and circumscribed around C . Then there are infinitely many of such n -gons.*




Figure 3: Pentagons inscribed and circumscribed

Proof. Let us consider one such n -gon and let us denote its sides by l_1, \dots, l_n . By hypothesis, there is a non-zero section $s \in H^0(E_{n,C})$ vanishing along the $\binom{n}{2}$ vertices $l_i \cap l_j$ (for $1 \leq i, j \leq n$ and $i \neq j$) of these lines. We denote by $Z(s)$ the set of these vertices.

Let us tensor the following exact sequence

$$0 \longrightarrow \mathcal{O}_{\mathbb{P}^2} \longrightarrow E_{n,C} \longrightarrow \mathcal{I}_{Z(s)}(n-1) \longrightarrow 0$$

by \mathcal{O}_D . Since $D \cap Z(s)$ consists of n -points, it induces the following decomposition of $E_{n,C}$ along D :

$$E_{n,C} \otimes \mathcal{O}_D = \mathcal{O}_D\left(\frac{n-2}{2}\right) \oplus \mathcal{O}_D\left(\frac{n}{2}\right).$$

According to this decomposition, we consider the following exact sequence:

$$0 \longrightarrow F \longrightarrow E_{n,C} \longrightarrow \mathcal{O}_D\left(\frac{n-2}{2}\right) \longrightarrow 0$$

where F is a rank two vector bundle over \mathbb{P}^2 . Taking the cohomology long exact sequence we verify immediately that $h^0(F) \geq 2$. Then, let us consider a pencil of sections of F and also the pencil of sections of $E_{n,C}$ induced by it. We obtain a commutative diagram:

$$\begin{array}{ccccccc}
& \mathcal{O}_{\mathbb{P}^2}^2 & \xlongequal{\quad} & \mathcal{O}_{\mathbb{P}^2}^2 & & & \\
& \downarrow & & \downarrow & & & \\
0 & \longrightarrow & F & \longrightarrow & E_{n,C} & \longrightarrow & \mathcal{O}_D(\frac{n-2}{2}) \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \parallel \\
0 & \longrightarrow & \mathcal{L}_1 & \longrightarrow & \mathcal{L}_2 & \longrightarrow & \mathcal{O}_D(\frac{n-2}{2}) \longrightarrow 0.
\end{array}$$

The sheaf \mathcal{L}_2 is supported by a curve Γ_2 of degree $(n-1)$ that is the determinant of a pencil of sections of $E_{n,C}$. This curve Γ_2 is a Poncelet curve. Then a general point on Γ_2 is a vertex of a complete n -gon inscribed in Γ_2 and circumscribed around C . Moreover any intersection point of the n lines forming the n -gon with Γ_2 is a vertex of this n -gon (it is clear by Bézout theorem since $n(n-1) = 2 \times \binom{n}{2}$). Let Γ_1 be the curve supporting the sheaf \mathcal{L}_1 ⁽³⁾. We have of course $\Gamma_2 = D \cup \Gamma_1$. Then D is an irreducible component of a Poncelet curve and by the way any general point on D is the vertex of complete n -gon inscribed in Γ_2 . Then this configuration meets the conic D in at least (because there are n lines) and at most (because they are vertices and the decomposition of the bundle along D is fixed) n points, so exactly n points, each counting twice. □

I thank the referee for the helpful remarks provided and for offering me the pictures 1 and 3.

References

- [1] W. Barth, Th. Bauer, Poncelet theorems, *Expo. Math.*, 14-2 (1996), 125–144.
- [2] M. Berger, *Géométrie vivante ou l'échelle de Jacob*, Cassini, 2009.
- [3] H.J.M. Bos, C. Kers, F. Oort, D.W. Raven, Poncelet's closure theorem, its history, its modern formulation, a comparison of its modern proof with those by Poncelet and Jacobi, and some mathematical remarks inspired by these early proofs, *Expo. Math.* 5 (1987), 289–364.

³As the referee pointed out to me, the existence of this curve can be seen directly, by computing the dimension of the vector space of polynomials of degree $(n-3)$ passing through the $\binom{n}{2} - n$ vertices that do not belong to D . This remark leads to a simplified version of this proof. The referee also reminded me that, according to Darboux ([4], livre III, chapitre II page 252), this curve is a union of $\frac{n-3}{2}$ conics for n odd, or a union of $\frac{n-4}{2}$ conics and a line for n even.

- [4] G. Darboux, *Principes de géométrie analytique*, Paris, Gauthier-Villars, 1917.
- [5] P. Griffiths, J. Harris, On Cayley's explicit solution to Poncelet's porism, *Enseign. Math.* (2) 24 (1978), no. 1-2, 31–40.
- [6] G. Ottaviani, E. Sernesi, On the hypersurface of Lüroth quartics, *Michigan Math. J.* 59 (2010), no. 2, 365–394.
- [7] J-V. Poncelet, *Traité des propriétés projectives des figures*, Bachelier Libraire, édition de 1822.
- [8] R. L. E. Schwarzenberger, Vector bundles on the projective plane, *Proc. London Math. Soc.* 11, (1961), 623–640.
- [9] J. Vallès, Fibrés de Schwarzenberger et coniques de droites sauteuses, *Bull. Soc. Math. France* 128, (2000), 433–449.